arXiv:1504.02918v2 [math.AP] 28 Sep 2015

GLOBAL REGULARITY CRITERION FOR THE 3D NAVIER-STOKES EQUATIONS WITH LARGE DATA

ABDELHAFID YOUNSI

Abstract. In this paper, we study the global regularity of strong solution to the Cauchy problem of 3D incompressible Navier-Stokes equations with large data and non-zero force. We prove that the strong solution exists globally for $\nabla u \in L^{\alpha}\left(0, T, L^{2}(\Omega)\right)$ with $2 \leq \alpha \leq 4$.

1. Introduction

Leray (1934) [8] and Hopf (1951) [6] showed the existence of global weak solutions to the three-dimensional Navier-Stokes system, but their uniqueness is still a fundamental open problem [4] and [5]. Furthermore, the strong solutions for the 3D Navier-Stokes equations are unique and can be shown to exist on a certain finite time interval for small initial data and small forcing term (see [2, 4, 5, 7] and references therein). The global existence of small strong solutions has been proved, see Constantin [4]. For the 3D Navier-Stokes equations with large data, we don't have a result of global existence and the question of asymptotic behavior of strong solution, as time variable t goes to $+\infty$ is a major problem. Most recently, there has been some progress along this line (see, for example, [2], [3] and [9]).

For $\nabla u \in L^{2}(0, T; L^{2}(\Omega))$, it known that for $\|\nabla u_{0}\|_{L^{2}}^{2}+\dfrac{2}{\nu} \int_{0}^{T}\|f\|_{L^{2}}^{2} \, ds \leq c \lambda_{1}^{1/2} \nu^{2}$ we have global regularity Constantin [4, P. 80 Theorem 9.3]. In the work of Beirao 1995 [1], it has been proved that the regularity is ensured if $\nabla u \in L^{4}(0, T; L^{2}(\Omega))$.

In this paper, we prove global regularity of strong solutions to the 3D Navier-Stokes problem in the classe $\nabla u \in L^{\alpha}\left(0, T, L^{2}(\Omega)\right)$ with $2 \leq \alpha \leq 4$. We give a sufficient condition for global existence in time for strong solution to the 3D Navier-Stokes equations with external force and large data.

2. The regularity criterion

In this paper, we consider the three-dimensional Navier-Stokes system

$$\begin{gathered}
\frac{\partial u}{\partial t}+u . \nabla u=\nu \triangle u+\nabla p+f, \ t>0, \\
\operatorname{div} u=0, \text { in } \Omega \times(0, \infty), \ u=0 \text { on } \partial \Omega \times(0, \infty) \text { and } u(x, 0)=u_{0}, \text { in } \Omega,
\end{gathered} \tag{2.1}$$

where $u=u(x, t)$ is the velocity vector field, $p(x, t)$ is a scalar pressure, f is a given force field and $\nu>0$ is the viscosity of the fluid. $u(x, 0)$ with $\operatorname{div} u_{0}=0$ in the sense of distribution is the initial velocity field, and Ω is a regular, open, bounded subset of $\mathbb{R}^{3}$ with smooth boundary $\partial \Omega$. We denote by $H^{m}(\Omega)$, the Sobolev space. We define the usual function spaces $\mathcal{V}=\left\{u \in C_{0}^{\infty}(\Omega): \operatorname{div} u=0\right\}$, $V=$ closure

---

2010 *Mathematics Subject Classification.* 76D05, 76D03.

Key words and phrases. Navier-Stokes equations. Strong solutions.

of $\mathcal{V}$ in $H_0^1(\Omega)$, $H =$ closure of $\mathcal{V}$ in $L^2(\Omega)$. The space H is equipped with the scalar product $(.,.)$ induced by $L^2(\Omega)$ and the norm $\|.\|_{L^2} = \|.\|_{L^2(\Omega)}$. For the local existence of strong solutions, we have the following result in 3D [4].

Theorem 2.1. [4, P. 82 Theorem 9.4] *Assume that* $u_0 \in V$ *and* $f \in L^2(0,T;H)$ *are given. Then there exists a* $T^* > 0$ *depending on* ν, f, u_0 *and* T, *such that there exists a unique solution of* (2.1) *satisfies* $u \in L^\infty(0,T^*;V) \cap L^2\left(0,T^*;H^2(\Omega)\right)$.

For $f = 0$, our main result on the global regularity for strong solutions with large data to the 3D Navier-Stokes equations (2.1) reads as follows

Theorem 2.2. *Let* $u_0 \in V$ *and* $u(x,t)$ *is the corresponding strong solution to the system* (2.1) *on* $[0,T]$. *Then if* u *satisfies*

$$\int_0^T \|\nabla u(.,t)\|_{L^2}^{\alpha}\, dt < C_1(\nu, u_0) \text{ for } 2 \le \alpha \le 4 \tag{2.2}$$

the strong solution $u(x,t)$ *exists globally in time for all finite value of* $\|\nabla u_0\|_{L^2}$.

Proof. Multiplying (2.1) by $\triangle u$ and integrate over Ω, we obtain

$$\frac{1}{2}\frac{d}{dt}\|\nabla u(.,t)\|_{L^2}^2 + \nu\|\triangle u\|_{L^2}^2 = ((u.\nabla u).\triangle u). \tag{2.3}$$

We will denote by $c_{i\in\mathbb{N}}$, positive constants. Using the Holder inequality and the Sobolev theorem, we get

$$|((u.\nabla u).\triangle u)| \le c_1 \|\nabla u\|_{L^2}^{\frac{3}{2}} \|\triangle u\|_{L^2}^{\frac{3}{2}}, \tag{2.4}$$

see [4, P. 79 (2.22)]. Combining (2.3) and (2.4), we obtain

$$\frac{d}{dt}\|\nabla u(.,t)\|_{L^2}^2 + \nu\|\triangle u\|_{L^2}^2 \le c_1 \|\nabla u\|_{L^2}^{\frac{3}{2}} \|\triangle u\|_{L^2}^{\frac{3}{2}}. \tag{2.5}$$

Setting $y = \|\nabla u(.,t)\|_{L^2}^2$ in (2.5), we get

$$y'(t) + \nu\|\triangle u\|_{L^2}^2 \le c_1 y^{\frac{3}{4}} \|\triangle u\|_{L^2}^{\frac{3}{2}}. \tag{2.6}$$

Dividing (2.6) by $(y+1)^2$ we get

$$\frac{1}{(y+1)^2}y'(t) + \nu\frac{\|\triangle u\|_{L^2}^2}{(y+1)^2} \le c_1 y^{\frac{3}{4}} \frac{\|\triangle u\|_{L^2}^{\frac{3}{2}}}{(y+1)^2}, \tag{2.7}$$

this is equivalant to

$$\frac{1}{(y+1)^2}y'(t) + \frac{\nu}{(y+1)^2}\|\triangle u\|_{L^2}^2 \le c_2 y^{\frac{\alpha}{8}} \frac{\|\triangle u\|_{L^2}^{\frac{3}{2}}}{(y+1)^{\frac{3}{2}}} \frac{y^{\frac{6-\alpha}{8}}}{(y+1)^{\frac{1}{2}}} \text{ with } 2 \le \alpha \le 4. \tag{2.8}$$

For $2 \le \alpha \le 4$ we have $\frac{y^{\frac{6-\alpha}{8}}}{(y+1)^{\frac{1}{2}}} \le 1$, then yields

$$\frac{1}{(y+1)^2}y'(t) + \frac{\nu}{(y+1)^2}\|\triangle u\|_{L^2}^2 \le c_3 y^{\frac{\alpha}{8}} \frac{\|\triangle u\|_{L^2}^{\frac{3}{2}}}{(y+1)^{\frac{3}{2}}}. \tag{2.9}$$

Applying Young's inequality on the second member of (2.9) with $p = 4/3$ and $q = 4$ we obtain

$$\frac{1}{(y+1)^2}y'(t) + \nu\frac{\|\triangle u\|_{L^2}^2}{(y+1)^2} \le c_4 \nu^3 y^{\frac{\alpha}{2}} + \nu\frac{\|\triangle u\|_{L^2}^2}{2(y+1)^2}, \tag{2.10}$$

wich gives

$$\frac{1}{(y+1)^2} y\prime(t) + \nu \frac{\|\triangle u\|_{L^2}^2}{2(y+1)^2} \leq c_5 \nu^3 y^{\frac{\alpha}{2}}. \tag{2.11}$$

If we drop the positive term $\nu \frac{\|\triangle u\|_{L^2}^2}{2(y+1)^2}$ in (2.11) then we have

$$\frac{1}{(y+1)^2} y\prime(t) \leq c_6 y^{\frac{\alpha}{2}} \text{ with } c_6 = c_5 \nu^3. \tag{2.12}$$

Multiplying (2.12) by $\sin\left(\frac{1}{y+1}\right)$ and using the fact that $\sin\left(\frac{1}{y+1}\right) \leq 1$, we obtain

$$\sin\left(\frac{1}{y+1}\right) \frac{1}{(y+1)^2} y\prime(t) \leq c_6 y^{\frac{\alpha}{2}}. \tag{2.13}$$

Integrating (2.13) on $[0,t]$, $t \leq T$, we get

$$\cos\left(\frac{1}{y+1}\right) \leq \cos\left(\frac{1}{y_0+1}\right) + c_6 \int_0^T y^{\frac{\alpha}{2}} dt. \tag{2.14}$$

For all finite y_0 we have $\cos\left(\frac{1}{y_0+1}\right) < 1$, then we can choose

$$\int_0^T y^{\frac{\alpha}{2}} dt < \frac{1 - \cos\left(\frac{1}{y_0+1}\right)}{2c_6} = C_1\left(\nu, \|\nabla u_0\|_{L^2}\right). \tag{2.15}$$

Thus, (2.15) yields $\cos\left(\frac{1}{y+1}\right) < 1$ and therefore $\frac{1}{y+1}$ can not identically equal to zero wich means that $y(t)$ is finite for all $t \geq 0$ and consequently $\nabla u(.,t) \in L^\infty\left(0,T;L^2(\Omega)\right)$. This proves, in particular that (2.2) is a condition sufficient for global regularity. □

For the 3D Navier-Stokes equations with no-negligible forces we prove the following result

Proposition 2.3. *Let $u_0 \in V$, $f \in L^2\left(0,T;L^2(\Omega)\right)$ and $u(x,t)$ is the corresponding strong solution to the system (2.1) on $[0,T]$. Then if u satisfies*

$$\int_0^T \|\nabla u(.,t)\|_{L^2}^\alpha dt < C_2(\nu, u_0, f) \text{ for } 2 \leq \alpha \leq 4 \tag{2.16}$$

the strong solution $u(x,t)$ exists globally in time for all finite value of $\|\nabla u_0\|_{L^2}$.

Proof. Using the Schwartz and Young inequalities, we obtain that

$$\begin{aligned} |(f, \triangle u)| &\leq \|f\|_{L^2} \|\triangle u\|_{L^2} \\ &\leq c_7 \|f\|_{L^2}^2 + \frac{\nu}{4} \|\triangle u\|_{L^2}^2. \end{aligned} \tag{2.17}$$

Combining (2.3), (2.4) and (2.17), we have

$$\frac{d}{dt} \|\nabla u(.,t)\|_{L^2}^2 + \nu \|\triangle u\|_{L^2}^2 \leq c_7 \|f\|_{L^2}^2 + c_1 y^{\frac{3}{4}} \|\triangle u\|_{L^2}^{\frac{3}{2}}. \tag{2.18}$$

Dividing (2.18) by $(y+1)^2$ and since $\frac{1}{(y+1)^2} \leq 1$, it follows that

$$\frac{1}{(y+1)^2} y\prime(t) + \nu \frac{\|\triangle u\|_{L^2}^2}{(y+1)^2} \leq c_7 \|f\|_{L^2}^2 + c_1 y^{\frac{3}{4}} \frac{\|\triangle u\|_{L^2}^{\frac{3}{2}}}{(y+1)^2}. \tag{2.19}$$

The proof of (2.16) follows in the same manner as that of (2.2). After some manipulations similaire to (2.3)-(2.13) and integration on $[0,T]$, we find

$$\cos\left(\frac{1}{y+1}\right) \leq \cos\left(\frac{1}{y_0+1}\right) + c_7\int_0^T \|f\|_{L^2}^2 + c_6\int_0^T y^{\frac{\alpha}{2}}, \tag{2.20}$$

which implies if

$$\cos\left(\frac{1}{y_0+1}\right) + c_7\int_0^T \|f\|_{L^2}^2 + c_6\int_0^T y^{\frac{\alpha}{2}} < 1, \tag{2.21}$$

then we have global regularity and $\nabla u\,(.,t) \in L^\infty\left(0,\infty; L^2\,(\Omega)\right)$. As before, we can choose

$$C_2\,(\nu,u_0,f) = \frac{1}{2c_6}\left(1 - \cos\left(\frac{1}{y_0+1}\right) - c_7\int_0^T \|f\|_{L^2}^2\right). \tag{2.22}$$

This finishes the proof of (2.16). □

References

[1] H. Beirao da Veiga, A New Regularity Class for the Navier-Stokes Equations in $\mathbb{R}^n$, Chin. Ann. of Math, 16B, 407 (1995).

[2] C. Cao, E. S. Titi, Regularity criteria for the three dimensional Navier-Stokes equations. Indiana Univ. Math. J. 57, 6, (2008) 2643-2661.

[3] J. Y. Chemin, I. Gallagher, M. Paicu, Global regularity for some classes of large solutions to the Navier-Stokes equations, Ann. of Math. (2) 173 , no. 2, (2011) 983-1012.

[4] P. Constantin, C. Foias, Navier-Stokes Equations. University of Chicago Press (1988).

[5] C. Foias, O. Manley, R. Rosa, R. Temam, Navier-Stokes Equations and Turbulence. Cambridge University Press. (2001).

[6] E. Hopf, Über die Anfangswertaufgabe für die hydrodynamischen Grun dgleichungen. Math. Nachr.4 (1951), 213-231.

[7] T. Kato, H. Fujita, On the nonstationary Navier-Stokes system., Rend. Sem. Mat. Univ. Padova 32, (1962) 243-260.

[8] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace. Acta Math. 63, (1934) 193-248.

[9] J. Xuanji, Z. Yong, Remarks on regularity criteria for the Navier-Stokes equations via one velocity component. Nonlinear Analysis: Real World Applications. 15, (2014) 239-245.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF DJELFA , ALGERIA.
E-mail address: `younsihafid@gmail.com`